

Rodanthi-Elisavet Mamouri^{1,2}, Argyro Nisantzi^{1,2}, George Kotsias¹, Hossein Panahifar¹, Maria Poutli^{1,2}, Konstantinos Chrysostomou^{1,2}, Athina Savva^{1,2}, Maria Prodromou^{1,2}, Georgia Charalambous^{1,2}, D. Hadjimitsis^{1,2}, Patric Seifert³ and Albert Ansmann³

1. Eratosthenes Centre of Excellence, Limassol, Cyprus
2. Cyprus University of Technology, Faculty of Engineering and Technology, Limassol, Cyprus
3. Leibniz Institut für Troposphärenforschung (TROPOS), Leipzig, Germany



A fresh smoke case

On July 23–24, 2025, a large wildfire originating near Mallia in the Limassol district of Cyprus generated an extensive smoke plume that affected air quality across the island and extended over the south-eastern Mediterranean.




The event was monitored in near real-time by the Cyprus Atmospheric Remote Sensing Observatory (CARO) National Facility (NF), in Limassol (LIM) [34.67° N, 33.04° E] operated by the ERATOSTHENES Centre of Excellence, using a suite of advanced ground-based instruments including PollyXT lidar, Doppler lidar, and cloud radar systems.

The July 2025 Limassol wildfire demonstrated in addition the value of EarthCARE's capabilities in active remote sensing of smoke plumes and underscored the strategic role of observatories like CARO-LIM in deepens scientific understanding of aerosol-radiation-cloud interactions, and in enhancing the scientific return and societal relevance of satellite missions.

The observatories are equipped with aerosol and cloud remote sensing instruments running continuously.

Table 1: Instruments deployed at LIM observatories for the EC VAL.

Limassol - CARO
PollyXT
AERONET phtometer
Wind lidar
Cloud Doppler Radar 35GHz
CHM15k ceilometer
Disdrometer
MWR HATPRO RPG
All sky camera
Pyranometer
Spectrophotometer



PlanetScope constellation : Burned area estimation (3m resolution)

The wildfire is estimated to have burned a total of 113.7 km². Taking into account that the total area of Cyprus is 9,251 km², the burned area corresponds to approximately 1.3% of the island's surface, a particularly significant percentage considering that it resulted from a single wildfire.

In addition, about 21% of the burned area falls within the Natura 2000 network, affecting important protected sites such as the Paramaliou River, the Cha Potami Special Protection Area, and the Limmati Valley Special Protection Area.

